

CSE4203: Computer Graphics  
Chapter – 6 (part - B)


# **Transformation Matrices**

Mohammad Imrul Jubair

# Outline

- 3D Linear Transformation
- 3D Scaling
- 3D Rotation
- Translation
- Affine Transformation

# Credit



## CS4620: Introduction to Computer Graphics

Cornell University

Instructor: Steve Marschner

<http://www.cs.cornell.edu/courses/cs4620/2019fa/>

# 3D Transformation (1/1)

- The linear 3D transforms are an extension of the 2D transforms.


- For 2D:

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} a_{11}x + a_{12}y \\ a_{21}x + a_{22}y \end{bmatrix}$$

- For 3D:

# 3D Scaling (1/1)

$$\text{scale}(s_x, s_y, s_z) = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & s_z \end{bmatrix}$$



# 3D Rotation (1/5)

- Rotation around axis
  - Counter-clockwise, w.r.t rotation axis.








Image Source: <https://slideplayer.com/slide/4889962/>





# 3D Rotation (2/5)

$$\text{rotate-z}(\phi) = \begin{bmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



# 3D Rotation (3/5)

$$\text{rotate-x}(\phi) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \phi & -\sin \phi \\ 0 & \sin \phi & \cos \phi \end{bmatrix}$$



# 3D Rotation (4/5)

$$\text{rotate-z}(\phi) = \begin{bmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{rotate-x}(\phi) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \phi & -\sin \phi \\ 0 & \sin \phi & \cos \phi \end{bmatrix}$$

$$\text{rotate-y}(\phi) = \begin{bmatrix} \cos \phi & 0 & \sin \phi \\ 0 & 1 & 0 \\ -\sin \phi & 0 & \cos \phi \end{bmatrix}$$

# 3D Rotation (5/5)

$$\text{rotate-z}(\phi) = \begin{bmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\text{rotate-x}(\phi) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \phi & -\sin \phi \\ 0 & \sin \phi & \cos \phi \end{bmatrix}$$


$$\text{rotate-y}(\phi) = \begin{bmatrix} \cos \phi & 0 & \sin \phi \\ 0 & 1 & 0 \\ -\sin \phi & 0 & \cos \phi \end{bmatrix}$$

Q: Why is it different?\*

\* <https://robotics.stackexchange.com/questions/10702/rotation-matrix-sign-convention-confusion>


# Translation in 2D (1/8)

- Move or Translate to another position.




# Translation in 2D (3/8)

original



translation



$$\begin{aligned}x' &= x + t_x \\y' &= y + t_y\end{aligned}$$

$$\mathbf{v}' = \mathbf{v} + \mathbf{t}$$

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} t_x \\ t_y \end{bmatrix} + \begin{bmatrix} x \\ y \end{bmatrix}$$



$$\mathbf{v}' = \mathbf{v} + \mathbf{t}$$

Source: <https://www.pling.org.uk/cs/cgv.html>

# Translation in 2D (4/8)

- But, for others cases, i.e. – scaling, rotation, we changed vectors  $\mathbf{v}$  using a **matrix  $M$** .
  - In 2D, these transforms have the form: -

$$\begin{aligned}x' &= m_{11}x + m_{12}y, \\y' &= m_{21}x + m_{22}y.\end{aligned}$$

$$\mathbf{v}' = \mathbf{M} \mathbf{v}$$

# Translation in 2D (5/8)

- We **cannot** use such transforms to **translate**, only to scale and rotate them.

$$\begin{aligned}x' &= m_{11}x + m_{12}y, \\y' &= m_{21}x + m_{22}y.\end{aligned}$$

$$\cancel{\mathbf{v}' = \mathbf{M} \mathbf{v}}$$





# Translation in 2D (6/8)

- There is just no way to do that by **multiplying**  $(x, y)$  by a  $2 \times 2$  matrix.
  - **adding** translation to our system of linear transformations:

$\begin{aligned}x' &= m_{11}x + m_{12}y, \\y' &= m_{21}x + m_{22}y.\end{aligned}$	$\mathbf{v}' = \mathbf{M} \mathbf{v}$
$\begin{aligned}x' &= x + x_t, \\y' &= y + y_t.\end{aligned}$	$\mathbf{v}' = \mathbf{v} + \mathbf{t}$

# Translation in 2D (7/8)

- This is perfectly feasible –



$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} t_x \\ t_y \end{bmatrix} + \begin{bmatrix} x \\ y \end{bmatrix}$$

$\underbrace{\hspace{10em}}$

$$\mathbf{v}' = \mathbf{v} + \mathbf{t}$$

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} s_x & 0 \\ 0 & s_y \end{bmatrix} \cdot \begin{bmatrix} x \\ y \end{bmatrix} \quad \begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix} \cdot \begin{bmatrix} x \\ y \end{bmatrix}$$

$\underbrace{\hspace{10em}}$

$$\mathbf{v}' = \mathbf{M} \mathbf{v}$$

Source: <https://www.pling.org.uk/cs/cgv.html>

# Translation in 2D (8/8)

- This is perfectly feasible
  - But, the rule for **composing transformations** is not as simple and clean as with linear transformations.

$$T = T_n \cdot T_{n-1} \dots T_1 \cdot T_0$$

$\begin{aligned}x' &= m_{11}x + m_{12}y, \\y' &= m_{21}x + m_{22}y.\end{aligned}$	$\mathbf{v}' = \mathbf{M} \mathbf{v}$
$\begin{aligned}x' &= x + x_t, \\y' &= y + y_t.\end{aligned}$	$\mathbf{v}' = \mathbf{v} + \mathbf{t}$

# Homogeneous Coordinates (1/9)

- Instead, we can use a clever trick to get a single matrix multiplication to do both.

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} & & \\ & 2 \times 2 & \\ & & \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

# Homogeneous Coordinates (2/9)

- Instead, we can use a clever trick to get a single matrix multiplication to do both.
- The idea is simple: represent the point  $(x, y)$  by a 3D vector  $[x \ y \ 1]^T$

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} & & \\ & 3 \times 3 & \\ & & \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

# Homogeneous Coordinates (3/9)

- Instead, we can use a clever trick to get a single matrix multiplication to do both.
- The idea is simple: represent the point  $(x, y)$  by a 3D vector  $[x \ y \ 1]^T$
- Use  $3 \times 3$  matrices of the form.

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} m_{11} & m_{12} & x_t \\ m_{21} & m_{22} & y_t \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

# Homogeneous Coordinates (4/9)

- This kind of transformation is called an *affine transformation*.
  - this way of implementing affine transformations by adding an extra dimension is called *homogeneous coordinates*

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} m_{11} & m_{12} & x_t \\ m_{21} & m_{22} & y_t \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

# Homogeneous Coordinates (5/9)

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} m_{11} & m_{12} & x_t \\ m_{21} & m_{22} & y_t \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

# Homogeneous Coordinates (6/9)

- Translation:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & x_t \\ 0 & 1 & y_t \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

# Homogeneous Coordinates (7/9)

- Scaling:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} S_x & 0 & 0 \\ 0 & S_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

# Homogeneous Coordinates (8/9)

- Rotation:

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

# 3D Transformation with Homogeneous Coordinates

## (1/1)

$$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$$

# 2D/ 3D Transformations (1/3)

	2D	3D
<b>T</b>	$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & x_t \\ 0 & 1 & y_t \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$	$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & p \\ 0 & 1 & 0 & q \\ 0 & 0 & 1 & r \\ 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$
<b>S</b>	$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} S_x & 0 & 0 \\ 0 & S_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$	$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} p & 0 & 0 & 0 \\ 0 & q & 0 & 0 \\ 0 & 0 & r & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$
<b>R</b>	$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos \phi & -\sin \phi & 0 \\ \sin \phi & \cos \phi & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$	$RotX = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos(\theta x) & -\sin(\theta x) & 0 \\ 0 & \sin(\theta x) & \cos(\theta x) & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ $RotY = \begin{bmatrix} \cos(\theta y) & 0 & \sin(\theta y) & 0 \\ 0 & 1 & 0 & 0 \\ -\sin(\theta y) & 0 & \cos(\theta y) & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ $RotZ = \begin{bmatrix} \cos(\theta z) & -\sin(\theta z) & 0 & 0 \\ \sin(\theta z) & \cos(\theta z) & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$

# Inverse Transformations (1/2)

<b>T</b>	<b>T<sup>-1</sup></b>
$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & p \\ 0 & 1 & 0 & q \\ 0 & 0 & 1 & r \\ 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$	$\begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & -p \\ 0 & 1 & 0 & -q \\ 0 & 0 & 1 & -r \\ 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix}$
$\begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix} = \begin{bmatrix} p & 0 & 0 & 0 \\ 0 & q & 0 & 0 \\ 0 & 0 & r & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$	$\begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} 1/p & 0 & 0 & 0 \\ 0 & 1/q & 0 & 0 \\ 0 & 0 & 1/r & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x' \\ y' \\ z' \\ 1 \end{bmatrix}$
$RotX = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos(\theta_x) & -\sin(\theta_x) & 0 \\ 0 & \sin(\theta_x) & \cos(\theta_x) & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ $RotY = \begin{bmatrix} \cos(\theta_y) & 0 & \sin(\theta_y) & 0 \\ 0 & 1 & 0 & 0 \\ -\sin(\theta_y) & 0 & \cos(\theta_y) & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ $RotZ = \begin{bmatrix} \cos(\theta_z) & -\sin(\theta_z) & 0 & 0 \\ \sin(\theta_z) & \cos(\theta_z) & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$	?




# Inverse Transformations (2/2)

	Transformation	Inverse Transformation
T	$T(tx, ty, tz)$	$T^{-1} = T(-tx, -ty, -tz)$
S	$S(sx, sy, sz)$	$S^{-1} = S(1/sx, 1/sy, 1/sz)$
R	$Rx(d)$ $Ry(d)$ $Rz(d)$	$R^{-1} = R(-d) = R^T$  $Rx^{-1} = Rx^T$ $Ry^{-1} = Ry^T$ $Rz^{-1} = Rz^T$

**Task:** take any transformation matrix (i.e. scaling matrix  $S$ ) with numerical values, do the matrix inversion and see if it becomes  $S^{-1}$


# Practice Problem - 1

- Scale w.r.t the center




# Practice Problem – 1 (Sol.)

- Scale w.r.t the center




# Practice Problem – 1 (Sol.)

- Scale w.r.t the center




# Practice Problem – 1 (Sol.)

- Scale w.r.t the center




# Practice Problem – 1 (Sol.)

- Scale w.r.t the center



# Practice Problem - 2

- We need to rotate a pyramid  $P$  about point  $(5, 5)$  by  $90^\circ$ . You have to –
  - Mention the steps to perform the task.
  - Determine the composite transformation matrix  $M$ .
  - Multiply  $M$  with  $P$  and determine the new coordinates  $P'$ .
  - Plot  $P$  and  $P'$  on the same axis to show the rotation.



# Additional Reading

- 3D Shearing
- 3D reflection
- Rigid-body transforms
- Windowing transformations

# Exercises

- Exercise 1 – 6, 8 and 9