


CSE4203: Computer Graphics  
Chapter – 8 (part - C)  
**Graphics Pipeline**

Mohammad Imrul Jubair

# Outline

- Barycentric Interpolation
- Rasterizing a triangle

# Credit




## CS4620: Introduction to Computer Graphics

Cornell University

Instructor: Steve Marschner


<http://www.cs.cornell.edu/courses/cs4620/2019fa/>

# Triangle Rasterization





Use Midpoint Algorithm for edges and fill in?


# Triangle Rasterization



# Triangle Rasterization




# Barycentric Coordinate



$$P(x, y) = \mathbf{O} + x\mathbf{X} + y\mathbf{Y}$$

$$P(\beta, \gamma) = a + \beta ? + \gamma ?$$

# Barycentric Coordinate



$$P(x, y) = \mathbf{O} + x\mathbf{X} + y\mathbf{Y}$$

$$P(\beta, \gamma) = a + \beta ? + \gamma ?$$


$$= a + \beta (b - a) + \gamma (c - a)$$

$$= a + \beta b - \beta a + \gamma c - \gamma a$$

$$= a + \beta b - \beta a + \gamma c - \gamma a$$



# Barycentric Coordinate



$$P(x, y) = \mathbf{O} + x\mathbf{X} + y\mathbf{Y}$$

$$P(\alpha, \beta, \gamma) = a + \beta ? + \gamma ?$$

$$= a + \beta (b - a) + \gamma (c - a)$$

$$= a + \beta b - \beta a + \gamma c - \gamma a$$

$$= a + \beta b - \beta a + \gamma c - \gamma a$$


$$= a(1 - \beta - \gamma) + b\beta + c\gamma$$

$$= \alpha a + \beta b + \gamma c$$

$$\begin{aligned} \alpha &= (1 - \beta - \gamma) \\ \Rightarrow \alpha + \beta + \gamma &= 1 \end{aligned}$$

In a barycentric coordinate system: **location of a point is specified by reference to a triangle for points in a plane.**

# Barycentric Coordinate



$$P(x, y) = \mathbf{O} + x\mathbf{X} + y\mathbf{Y}$$

$$P(\alpha, \beta, \gamma) = \alpha\mathbf{a} + \beta\mathbf{b} + \gamma\mathbf{c}$$

Barycentric


$$\alpha + \beta + \gamma = 1$$

$$0 < \alpha < 1,$$

$$0 < \beta < 1,$$

$$0 < \gamma < 1.$$

# Barycentric Coordinate



$$P(x, y) = \mathbf{O} + x\mathbf{X} + y\mathbf{Y}$$

$$P(\alpha, \beta, \gamma) = \alpha\mathbf{a} + \beta\mathbf{b} + \gamma\mathbf{c}$$

$$\alpha + \beta + \gamma = 1$$

$$0 < \alpha < 1,$$


$$0 < \beta < 1,$$

$$0 < \gamma < 1.$$

*Cartesian*  $\rightarrow$  *Barycentric*

$P(x, y) \rightarrow P(\alpha, \beta, \gamma)$

# Cartesian $\rightarrow$ Barycentric



$$y = mx + b$$

$$\Rightarrow y - mx - b = 0$$


$$\Rightarrow y - \frac{y_c - y_a}{x_c - x_a}x - b = 0$$

$$\Rightarrow (x_c - x_a)y - (y_c - y_a)x - (x_c - x_a)b = 0$$

$$\Rightarrow (x_c - x_a)y - (y_c - y_a)x - C = 0$$


$$\Rightarrow (y_a - y_c)x + (x_c - x_a)y - C = 0$$

# Cartesian $\rightarrow$ Barycentric



$$\begin{aligned} C &= (y_a - y_c)x + (x_c - x_a)y \\ &= (y_a - y_c)\mathbf{x}_a + (x_c - x_a)\mathbf{y}_a \\ &= x_c y_a - x_a y_c \end{aligned}$$


# Cartesian $\rightarrow$ Barycentric




$$\begin{aligned} C &= (y_a - y_c)x + (x_c - x_a)y \\ &= (y_a - y_c)x_a + (x_c - x_a)y_a \\ &= x_c y_a - x_a y_c \end{aligned}$$

$$f_{ac}(x, y) = (y_a - y_c)x + (x_c - x_a)y - x_c y_a + x_a y_c = 0$$

# Cartesian $\rightarrow$ Barycentric




# Cartesian $\rightarrow$ Barycentric



$$\beta = \frac{f_{ac}(x, y)}{f_{ac}(x_b, y_b)}$$



# Cartesian $\rightarrow$ Barycentric




$$\beta = \frac{f_{ac}(x, y)}{f_{ac}(x_b, y_b)}$$

$$= \frac{(y_a - y_c)x + (x_c - x_a)y + x_a y_c - x_c y_a}{(y_a - y_c)x_b + (x_c - x_a)y_b + x_a y_c - x_c y_a},$$

**Question – 1:** In which case  $\beta$  becomes 1?

**Question – 2:** What will happen when  $(x, y)$  lies on  $f_{ab}(x, y)$


# Cartesian $\rightarrow$ Barycentric



$$\beta = \frac{f_{ac}(x, y)}{f_{ac}(x_b, y_b)}$$

$$\gamma = \frac{f_{ac}(x, y)}{f_{ac}(x_b, y_b)}$$

# Cartesian $\rightarrow$ Barycentric




$$\beta = \frac{f_{ac}(x, y)}{f_{ac}(x_b, y_b)}$$

$$\gamma = \frac{f_{ac}(x, y)}{f_{ac}(x_b, y_b)}$$

$$\alpha = 1 - \beta - \gamma$$

# Cartesian $\rightarrow$ Barycentric



$$P(x, y) \rightarrow P(\alpha, \beta, \gamma)$$


$$\beta = \frac{f_{ac}(x, y)}{f_{ac}(x_b, y_b)}$$

$$\gamma = \frac{f_{ac}(x, y)}{f_{ac}(x_b, y_b)}$$

$$\alpha = 1 - \beta - \gamma$$


# Barycentric Coordinate

Another approach:




Credit: Fundamentals of Computer Graphics 3<sup>rd</sup> Edition by Peter Shirley, Steve Marschner | <http://www.cs.cornell.edu/courses/cs4620/2019fa/>

# Triangle Rasterization (1/7)



Use Midpoint Algorithm for edges and fill in?




Use an approach based on barycentric coordinates

# Triangle Rasterization (2/7)

- If the vertices have colors  $c_0$ ,  $c_1$ , and  $c_2$ , the color at a point in the triangle with *Barycentric coordinates*  $(\alpha, \beta, \gamma)$  is:


$$\mathbf{c} = \alpha\mathbf{c}_0 + \beta\mathbf{c}_1 + \gamma\mathbf{c}_2$$



- This type of interpolation of color is known in graphics as ***Gouraud interpolation***

# Triangle Rasterization (3/7)


```
for all  $x$  do  
  for all  $y$  do  
    compute  $(\alpha, \beta, \gamma)$  for  $(x, y)$   
    if  $(\alpha \in [0, 1]$  and  $\beta \in [0, 1]$  and  $\gamma \in [0, 1])$  then  
       $\mathbf{c} = \alpha\mathbf{c}_0 + \beta\mathbf{c}_1 + \gamma\mathbf{c}_2$   
      drawpixel  $(x, y)$  with color  $\mathbf{c}$ 
```






# Triangle Rasterization (4/7)

```
for  $y = y_{\min}$  to  $y_{\max}$  do  
  for  $x = x_{\min}$  to  $x_{\max}$  do  
  
    compute  $(\alpha, \beta, \gamma)$  for  $(x, y)$   
  
    if  $(\alpha > 0$  and  $\beta > 0$  and  $\gamma > 0)$  then  
       $\mathbf{c} = \alpha \mathbf{c}_0 + \beta \mathbf{c}_1 + \gamma \mathbf{c}_2$   
      drawpixel  $(x, y)$  with color  $\mathbf{c}$ 
```



# Triangle Rasterization (5/7)

```
for  $y = y_{\min}$  to  $y_{\max}$  do  
  for  $x = x_{\min}$  to  $x_{\max}$  do  
     $\alpha = f_{12}(x, y) / f_{12}(x_0, y_0)$   
     $\beta = f_{20}(x, y) / f_{20}(x_1, y_1)$   
     $\gamma = f_{01}(x, y) / f_{01}(x_2, y_2)$   
    if ( $\alpha > 0$  and  $\beta > 0$  and  $\gamma > 0$ ) then  
       $\mathbf{c} = \alpha \mathbf{c}_0 + \beta \mathbf{c}_1 + \gamma \mathbf{c}_2$   
      drawpixel ( $x, y$ ) with color  $\mathbf{c}$ 
```



# Triangle Rasterization (6/7)

**for**  $y = y_{\min}$  **to**  $y_{\max}$  **do**

**for**  $x = x_{\min}$  **to**  $x_{\max}$  **do**

$$\alpha = f_{12}(x, y) / f_{12}(x_0, y_0)$$

$$\beta = f_{20}(x, y) / f_{20}(x_1, y_1)$$


$$\gamma = f_{01}(x, y) / f_{01}(x_2, y_2)$$

**if** ( $\alpha > 0$  and  $\beta > 0$  and  $\gamma > 0$ ) **then**


$$\mathbf{c} = \alpha \mathbf{c}_0 + \beta \mathbf{c}_1 + \gamma \mathbf{c}_2$$

    drawpixel  $(x, y)$  with color  $\mathbf{c}$

$$f_{01}(x, y) = (y_0 - y_1)x + (x_1 - x_0)y + x_0y_1 - x_1y_0,$$
$$f_{12}(x, y) = (y_1 - y_2)x + (x_2 - x_1)y + x_1y_2 - x_2y_1,$$
$$f_{20}(x, y) = (y_2 - y_0)x + (x_0 - x_2)y + x_2y_0 - x_0y_2.$$



# Triangle Rasterization (7/7)



					0.00	
					1.00	
					0.00	
				0.25	0.25	
				0.75	1.00	
				0.00	0.00	
			0.50	0.50	0.50	
			0.50	0.75	1.00	
			0.00	0.00	0.00	
		0.75	0.75	0.75	0.75	
		0.25	0.50	0.75	1.00	
		0.00	0.00	0.00	0.00	
	1.00	1.00	1.00	1.00	1.00	
	0.00	0.25	0.50	0.75	1.00	
	0.00	0.00	0.00	0.00	0.00	

Credit: Fundamentals of Computer Graphics 3<sup>rd</sup> Edition by Peter Shirley, Steve Marschner | <http://www.cs.cornell.edu/courses/cs4620/2019fa/>

# Practice Problem

- Take three vertices of a triangle, choose two points, P and Q , such that they stay inside and outside the triangle respectively.
  - Apply barycentric interpolation and verify that P lies inside and Q lies outside the triangle.